

PATIENT _____

DATE	PROBLEM	GOAL	TO DATE	INTERVENTIONS	RESP DISC
	<p>Heart Failure</p> <p>related to</p> <p>Coronary Artery Disease</p> <p>Cardiomyopathy</p> <p>Congenital Heart Disease</p> <p>Heart Valve disease</p> <p>Arrythmia</p> <p>Patient has exhibited symptoms indicative of heart failure since the previous assessment.</p> <p>Actions taken to respond:</p> <p>Patient's physician contacted the same day</p> <p>Patient advised to get emergency treatment</p> <p>Implement physician-ordered patient-specific established parameters for treatment</p> <p>Patient education or other clinical interventions</p> <p>Obtained change in care plan orders (increased monitoring by agency, change in visit frequency, telehealth, etc.)</p>	<p>Patient safety will be maintained, and patient will be free of discomfort</p> <p>Patient / caregivers will verbalize understanding of diagnosis and treatment</p> <p>Patient / caregivers will demonstrate correct administration of medications</p>		<p>Monitor blood pressure</p> <p>Monitor for shortness of breath, weakness, dizziness, chest pain, jugular vein distention, decreased peripheral pulses, arrhythmias, fatigue, rales, dyspnea, changes in mental status, cool clammy skin, edema, cough, weight gain, fatigue, loss of appetite</p> <p>Monitor respiratory rate and depth, lung sounds</p> <p>Labs as ordered</p> <p>Instruct patient / caregivers to:</p> <p>Monitor intake and output</p> <p>Monitor weekly weight</p> <p>Monitor urine color</p> <p>Administer oxygen as ordered</p> <p>Alternate activities with periods of rest</p> <p>Administer medications and treatments as ordered</p> <p>Avoid straining at stool</p> <p>Limit salt intake</p>	